

High-latitude Input for Meso-scale Electrodynamics: HIME

Dogacan Su Ozturk¹

Xing Meng¹, Olga Verkhoglyadova¹,

Roger Varney², Ashton Reimer², Josh Semeter³

1: Jet Propulsion Laboratory, California Institute of Technology:

2: Center for Geospace Studies, SRI International, Menlo Park

3: Center for Space Physics, Boston University



Jet Propulsion Laboratory
California Institute of Technology

Ionospheric Structuring: In Situ and Groundbased Low Altitude Studies (ISINGLASS) Experiment

PI: Kristina Lynch – Dartmouth College



Robert Clayton (Dartmouth), Matt Zettergren (ERAU), Meghan Burleigh (ERAU-UMich), Mark Conde (UAF), Guy Grubbs (GSFC), Don Hampton (UAF), David Hysell (Cornell), Marc Lessard (UNH), Robert Michell (UMD), Ashton Reimer (SRI), T. Maximillian Roberts (Dartmouth-JPL), Marilia Samara (GSFC), Roger Varney (SRI)

Aim: Sampling multiple locations simultaneously in the auroral ionosphere to take gradient measurements of plasma parameters.

[Clayton et al., 2019a, b]



Running a global I-T model with ISR estimates of electric field



Modelling:

- PFISR aiding the ISINGLASS experiment
- The electric field estimates from 15 beams:
 - Temporal resolution [66 seconds]
 - Spatial resolution [0.05° in lat and 0.3° in lon, ~ 10 km]
 - Down-sampled to $0.75^\circ \times 0.75^\circ$: 35x80 km


Validation:

- Plasma measurements along Beams



High-latitude Input for Meso-scale Electrodynamics: HIME







- Nicolls et al. (2014) technique estimates F-region electric fields on a 2-D grid from the PFISR V_{LOS} measurements.
- These estimates can be merged with a global empirical potential model to drive global I-T models and investigate the role of meso-scale electric fields.



Methodology IV: Simulation setup

E fields at 20170302_0639

Subtracted Average = 30 minutes




- Weimer model results are weaker (~0-10 mV/m) than the PFISR electric field values.
- To assess the performance of the new model vs understanding the effects of variability:

$$E_{\text{total}} = E_{\text{background}} + E_{\text{variability}}$$

where $E_{\text{background}}$ 30 min. boxcar average.

Simulations	Potentials
1	Weimer Electric Field Potentials
2	Measured Electric Field Potentials
3	Background Electric Field Potentials
4	Variable Electric Field Potentials

Running a global I-T model with HIME




HIME is an add-on framework and does not require major source code modification.

Global Ionosphere -Thermosphere Model (GITM)

- 3D, altitude based non-uniform grid, assumes non-hydrostatic solution
- High-latitude input: Electric potential and particle precipitation
- Output: Plasma and neutral density, temperature, ion and electron velocity, neutral winds

Ridley, Deng and Toth, JASTP, 2006


Meso-scale electric fields in GITM



- ■ ■ PFISR Estimates
- Errors

* Profiles extracted from upper boundary.


Meso-scale electric fields in GITM



* Profiles extracted from upper boundary.

- ■ ■ PFISR Estimates
- Errors
- Weimer-driven GITM Runs


Meso-scale electric fields in GITM



- ■ ■ PFISR Estimates
- Errors
- Weimer-driven GITM Runs
- HIME[B]-driven GITM Runs

* Profiles extracted from upper boundary.


Meso-scale electric fields in GITM



- ■ ■ PFISR Estimates
- Errors
- Weimer-driven GITM Runs
- HIME[B]-driven GITM Runs
- HIME[V]-driven GITM Runs

* Profiles extracted from upper boundary.

Meso-scale electric fields in GITM




* Profiles extracted from upper boundary.

- ■ ■ PFISR Estimates
- Errors
- Weimer-driven GITM Runs
- HIME[B]-driven GITM Runs
- HIME[V]-driven GITM Runs
- HIME[T]-driven GITM Runs

- Both variability and amplitude are better captured with HIME.
- Electric field estimates from PFISR successfully incorporated.

Energy transfer rate with meso-scale electric fields



- Meso- and small- scale structures have been shown to have a profound effect on local heating*.
- The locally deposited energy increases in HIME-driven simulation compared to Weimer-driven simulation.

* Verkhoglyadova et al., 2018;
Lotko and Zhang, 2019

Effects of Meso-scale electric fields in the I-T system

For the first time a framework has been developed to incorporate ISR estimates of 2D electric fields to global I-T models.

The HIME driven simulations showed that meso-scale electric fields can lead to:

- Various changes in plasma parameters.
- Up to an order of magnitude increase in volumetric heating along the altitudinal profiles.

Ongoing and future work on HIME:

- A self-consistent treatment of particle precipitation to fully understand the effects of meso-scale structures.

Thank you.

Friday 13th December 2019, 15:28-15:40

SA53A-09: Results from a new approach to meso-scale driving of the I-T system

Acknowledgements

- This work is funded by the NASA ROSES 2016 Heliophysics LWS Science (NRA NNH16ZDA001N) Program.
- This work was performed at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.
- This material is based upon work supported by the Poker Flat Incoherent Scatter Radar which is a major facility funded by the National Science Foundation through cooperative agreement AGS-1840962 to SRI International.
- GITM is developed and supported by Prof. Aaron Ridley at University of Michigan.
- Simulations were done on NASA High-End Computing Program through the NASA Advanced Supercomputing Division at Ames Research Center, and Stampede Supercomputer at Texas Advanced Computing Center at University of Texas at Austin.
- We acknowledge use of NASA/GSFC's Space Physics Data Facility's OMNIWeb (or CDAWeb or ftp) service, and OMNI data.
- Part of the simulation results have been provided by the Community Coordinated Modeling Center at Goddard Space Flight Center through their public Runs on Request system (<http://ccmc.gsfc.nasa.gov>). The Weimer Model was developed by Daniel R. Weimer at Virginia Tech. The Ovation Prime Model was developed by Patrick Newell at JHU/APL.
- The authors gratefully acknowledge Mark Conde for maintaining the SDI data at [\url{https://sdi_server.gi.alaska.edu/sdiweb/index.asp}](https://sdi_server.gi.alaska.edu/sdiweb/index.asp) and ASI data at [\url{https://optics.gi.alaska.edu/optics/}](https://optics.gi.alaska.edu/optics/).
- We thank the developers of Spacepy ([\url{https://github.com/spacepy/spacepy.git}](https://github.com/spacepy/spacepy.git)), AACGMV2 ([\url{https://github.com/aburrell/aacgm2.git}](https://github.com/aburrell/aacgm2.git)), and Dascutils ([\url{https://github.com/space-physics/dascutils}](https://github.com/space-physics/dascutils)) libraries, which were used in this study.



Jet Propulsion Laboratory
California Institute of Technology

jpl.nasa.gov